

NEW TWISTED QUANTUM CURRENT ALGEBRAS

NAIHUAN JING

ABSTRACT. We introduce a twisted quantum affine algebra associated to each simply laced finite dimensional simple Lie algebra. This new algebra is a Hopf algebra with a Drinfeld-type comultiplication. We obtain this algebra by considering its vertex representation. The vertex representation quantizes the twisted vertex operators of Lepowsky-Wilson and Frenkel-Lepowsky-Meurman. We also introduce a twisted quantum loop algebra for the Kac-Moody case and give its level one representation.

1. INTRODUCTION

Let $\mathfrak{g}$ be the complex finite dimensional simple Lie algebra of simply laced type. Let $\alpha_1, \dots, \alpha_l$ be the fixed set of simple roots. The associated standard form is then defined by

$$(1.1) \quad (\alpha_i | \alpha_j) = a_{ij},$$

where $A = (a_{ij})$ is the Cartan matrix.

Let q be a generic complex number (i.e. not a root of unity). We define the q -integer $[n]$ by

$$[n] = \frac{q^n - q^{-n}}{q - q^{-1}}.$$

Other q -numbers in this paper are defined accordingly using the same base. Thus the q -factorial $[n]!$ is defined by $[n]! = \prod_{i=1}^n [i]$.

In the theory of quantum affine algebras [Dr, FJ] the q -deformed Heisenberg algebra $U_q(\widehat{\mathfrak{h}})$ inside $U_q(\widehat{\mathfrak{g}})$ is an associative algebra generated by $a_i(m)$ ($m \in \mathbb{Z}^\times$) and the central element γ subject to the following relations

$$(1.2) \quad [a_i(m), a_j(n)] = \delta_{m,-n} \frac{[(\alpha_i | \alpha_j)m]}{m} \frac{\gamma^m - \gamma^{-m}}{q - q^{-1}},$$

$$(1.3) \quad [a_i(m), \gamma] = 0.$$

In [FJ] we used this q -deformed Heisenberg algebra to give the first explicit construction of level one irreducible modules of the quantum affine algebras of simply laced types. Later the author [J2] generalized this construction to the twisted quantum affine algebras using some twisted q -Heisenberg algebras. Recently in [J7] we introduced a quantum loop algebra for the quantum Kac-Moody algebra by considering the bilinear form as any symmetric form on an even lattice.

Date: January 15, 1999; revised on April 27, 1999.

1991 Mathematics Subject Classification. Primary: 17B.

Key words and phrases. vertex operators, quantum affine algebras.

Research supported in part by NSA grant MDA904-97-1-0062 and NSF grant DMS-9701755 at MSRI.

The purpose of this paper is to study an analogous Heisenberg algebra and obtain a new twisted quantum affine algebra. Following the familiar twisting in conformal field theory [FLM, L] we will restrict the index set to be the set of odd integers in (1.2). With the help of this twisted Heisenberg algebra we will construct a new class of twisted quantum algebras and study their properties. Our construction also follows the more general twisting discussed in [J4].

The classical twisted vertex operators played a special role in the construction of the moonshine module $V^\natural$ [FLM] for the Monster group. It also helped build the twisted boson-fermion correspondence for the double covering group of S_n [J3]. One can also trace its appearance in various other realizations of affine Lie algebras [LP, DL] and their quantum analogs [J5, DF].

Part of our motivation is to construct a principal bosonization of the quantum affine algebra $U_q(\widehat{sl}_2)$, which seems quite non-trivial and is still unresolved. The new class of twisted quantum algebras constructed in this paper is different from that of the twisted quantum affine algebras given in [J2]. It is a common phenomenon that one classical object may have different q -deformations. It is unclear what will be the relation between our new algebras and the untwisted quantum affine algebras, though at least in the case of $\widehat{sl}_2$ the author conjectures that there exists a homomorphism from $U_q(\widehat{sl}_2)$ into the endomorphism ring of the Fock space constructed in section 2. We suggest that the verification of this conjecture would provide a principal construction for $U_q(\widehat{sl}_2)$. The difficulty seems to be the lack of a Drinfeld-Jimbo presentation for our twisted quantum current algebra.

Replacing the underlying root lattice by the root lattice of a Kac-Moody algebra, we also introduce a twisted quantum loop algebra associated to any symmetrizable generalized Cartan matrix. The vertex operator representation of this twisted Kac-Moody loop algebra uses some q -deformation of the rational function $(\frac{1-x}{1+x})^r$. The commutation relations in this case involve q -differences of the delta function.

The twisted current algebras first appeared in our work [J1] for the simplest case of A_1 . The algebra is similar to certain cases of Ding and Iohara's algebras [DI], though we have relatively stricter relations. It appears that a certain quotient of our algebra is also a quotient of their algebra. One interesting phenomenon (see [DI]) is that the Serre relations involve with non-constant coefficients. In our case we replace the Serre relations by commutation relations between two operators of same type (see (3.12-3.13)) as in the classical case [FLM]. I thank Jintai Ding for helpful discussions and comments on the paper.

2. TWISTED VERTEX REPRESENTATIONS

Let Q be the root lattice of the simple finite dimensional Lie algebra $\mathfrak{g}$ of the simply laced type with the standard bilinear form $(\cdot | \cdot)$ such that $(\alpha_i | \alpha_j) = a_{ij}$.

Let the q -deformed twisted Heisenberg algebra $U_q(\tilde{\mathfrak{h}})$ be the associative algebra generated by $a_i(m)$ ($m \in 2\mathbb{Z} + 1$) and the central element γ subject to the following relations

$$(2.1) \quad [a_i(m), a_j(n)] = \delta_{m, -n} \frac{[(\alpha_i | \alpha_j)m]}{2m} \frac{\gamma^m - \gamma^{-m}}{q - q^{-1}},$$

$$(2.2) \quad [a_i(m), \gamma] = [a_i(m), \gamma^{-1}] = 0.$$

The algebra $U_q(\tilde{\mathfrak{h}})$ has a canonical representation realized on $V = S(\tilde{\mathfrak{h}}^-)$, the space of symmetric polynomials in $a_i(-n)$, $n \in 2\mathbb{Z} + 1$, $n > 0$. This representation is

actually the induced one from the trivial representation $1_{\mathbb{C}}$ with the central charge $\gamma = q$ of the subalgebra $U_q(\mathfrak{h}^+)$ generated by $\gamma^{\pm 1}$ and $a_i(n), n \in 2\mathbb{Z} + 1, n > 0$. The explicit action is given by

$$(2.3) \quad \begin{aligned} \gamma^{\pm 1} &= \text{the multiplication operator by } q^{\pm 1}, \\ a_i(-n) &= \text{the multiplication operator by } a_i(-n), \\ a_i(n) &= \text{the annihilation operator by } a_i(n) \text{ subject to (2.1),} \end{aligned}$$

for $n \in 2\mathbb{Z} + 1, n > 0$.

We also denote by $a_i(n)$ be the operator corresponding to $a_i(n)$, then they satisfy the relation (2.1).

Let $\hat{Q}$ be the central extension of the root lattice Q such that

$$(2.4) \quad 1 \longrightarrow \mathbb{Z}_2 \longrightarrow \hat{Q} \longrightarrow Q \longrightarrow 1$$

with the commutator map

$$(2.5) \quad aba^{-1}b^{-1} = (-1)^{(\alpha|\beta)},$$

where we let a be the preimage of α . In the following we will denote the preimage of α_i by a_i . Let T be any $\hat{Q}$ -module such that as operators on T

$$(2.6) \quad a_i a_j = (-1)^{(\alpha_i|\alpha_j)} a_j a_i.$$

Let $V_Q = S(\tilde{\mathfrak{h}}^-) \otimes T$. We introduce the twisted vertex operators acting on V_Q by the following expressions:

$$(2.7) \quad E_-^{\pm}(\alpha_i, z) = \exp\left(\pm \sum_{n=1, \text{odd}}^{\infty} \frac{2q^{\mp n/2}}{[n]} a_i(-n) z^n\right),$$

$$(2.8) \quad E_+^{\pm}(\alpha_i, z) = \exp\left(\mp \sum_{n=1, \text{odd}}^{\infty} \frac{2q^{\mp n/2}}{[n]} a_i(n) z^{-n}\right),$$

$$(2.9) \quad X_i^{\pm}(z) = E_-^{\pm}(\alpha_i, z) E_+^{\pm}(\alpha_i, z) a_i^{\pm 1} = \sum_{n \in \mathbb{Z}} X_i^{\pm}(n) z^{-n}.$$

We define the normal ordering as usual by moving the annihilation operators to the right of the creation operators. It is a routine calculation to obtain the operator product expansions for the twisted vertex operators.

Proposition 2.1.

$$\begin{aligned} X_i^{\pm}(z) X_j^{\pm}(w) &=: X_i^{\pm}(z) X_j^{\pm}(w) : \begin{cases} 1 & \text{if } (\alpha_i|\alpha_j) = 0 \\ \frac{z-w}{z+w} \frac{z-q^{\pm 2}w}{z+q^{\pm 2}w} & \text{if } i = j \\ \frac{z+q^{\mp 1}w}{z-q^{\mp 1}w} & \text{if } (\alpha_i|\alpha_j) = -1 \end{cases} \\ X_i^{\pm}(z) X_j^{\mp}(w) &=: X_i^{\pm}(z) X_j^{\mp}(w) : \begin{cases} 1 & \text{if } (\alpha_i|\alpha_j) = 0 \\ \frac{z+qw}{z-qw} \frac{z+q^{-1}w}{z-q^{-1}w} & \text{if } i = j \\ \frac{z-w}{z+w} & \text{if } (\alpha_i|\alpha_j) = -1 \end{cases} \end{aligned}$$

In particular we have the following useful relations.

Lemma 2.1. *The operators $\Phi_i(zq^{-1/2}) =: X_i^+(zq^{-1})X_i^-(z)$: and $\Psi_i(zq^{1/2}) =: X_i^+(zq)X_i^-(z)$: are given by*

$$(2.10) \quad \Phi_i(z) = \exp((q^{-1} - q) \sum_{n>0, \text{odd}} a_i(-n)z^n) = \sum_{n \geq 0} \Phi_i(-n)z^n,$$

$$(2.11) \quad \Psi_i(z) = \exp((q - q^{-1}) \sum_{n>0, \text{odd}} a_i(n)z^{-n}) = \sum_{n \geq 0} \Psi_i(n)z^{-n}.$$

These two exponential operators can generate a complete system of basis for the q -Heisenberg algebra [FJ, J2]. Let $G_{ij}(z) = \sum_{n \geq 0} G_n z^n$ be the Taylor series of the function

$$(2.12) \quad \frac{q^{(\alpha_i|\alpha_j)}z - 1}{q^{(\alpha_i|\alpha_j)}z + 1} \frac{z + q^{(\alpha_i|\alpha_j)}}{z - q^{(\alpha_i|\alpha_j)}}$$

at $z = 0$.

Lemma 2.2. *The relations (1.5) for the Heisenberg algebras are equivalent to the following formal series one:*

$$\Phi_i(z)\Psi_j(w) = \Psi_j(w)\Phi_i(z)G_{ij}\left(\frac{z}{w}q^{-1}\right)/G_{ij}\left(\frac{z}{w}q\right).$$

Proof. From (1.5) it follows that

$$\begin{aligned} & \Phi_i(z)\Psi_j(w) \\ &= \Psi_j(w)\Phi_i(z) \exp \left\{ \sum_{n \geq 0, \text{odd}} \frac{2}{n} (q^{(\alpha_i|\alpha_j)n} - q^{-(\alpha_i|\alpha_j)n})(q^n - q^{-n}) \left(\frac{z}{w}\right)^n \right\} \\ &= \Psi_j(w)\Phi_i(z)G_{ij}(z/qw)/G_{ij}(qz/w), \end{aligned}$$

where we use the definition of $G_{ij}(z)$. □

It is easy to see that Lemma (2.2) is equivalent to the Heisenberg commutation relations (2.1). We give the complete relations satisfied by the twisted q -vertex operators as follows.

Theorem 2.3. *The twisted vertex operators obey the relations in Lemma 2.2 and the following relations:*

$$(2.13) \quad [\Phi_i(z), \Phi_j(w)] = [\Psi_i(z), \Psi_j(w)] = 0,$$

$$(2.14) \quad \Phi_i(z)X_j^\pm(w)\Phi_i(z)^{-1} = X_j^\pm(w)G_{ij}(q^{\mp 1/2}z/w)^{\pm 1},$$

$$(2.15) \quad \Psi_i(z)X_j^\pm(w)\Psi_i(z)^{-1} = X_j^\pm(w)G_{ij}(q^{\mp 1/2}w/z)^{\pm 1},$$

$$(2.16) \quad [X_i^\pm(z), X_j^\pm(w)] = [X_i^+(z), X_j^-(w)] = 0, \quad (\alpha_i|\alpha_j) = 0$$

$$(2.17) \quad [X_i^+(z), X_j^-(w)] = 2z : X_i^+(z)X_j^-(w) : \delta(-\frac{w}{z}), \quad (\alpha_i|\alpha_j) = -1$$

$$(2.18) \quad [X_i^+(z), X_i^-(w)] = \frac{2(q+q^{-1})}{q-q^{-1}} \left(\Psi_i(q^{-1/2}z)\delta(\frac{w}{z}q) - \Phi_i(zq^{1/2})\delta(\frac{w}{z}q^{-1}) \right),$$

$$(z - q^{\mp 1}w)(z + q^{\pm 1}w)X_i^\pm(z)X_j^\pm(w) =$$

$$(q^{\mp 1}z - w)(q^{\pm 1}z + w)X_j^\pm(w)X_i^\pm(z), \quad (\alpha_i|\alpha_j) = -1,$$

$$(2.19) \quad (z - q^{\pm 2}w)(z + q^{\mp 2}w)X_i^\pm(z)X_i^\pm(w) = (q^{\pm 2}z - w)(q^{\mp 2}z + w)X_i^\pm(w)X_i^\pm(z)$$

$$(2.20) \quad + 2(q + q^{-1})^2 \delta(-\frac{w}{z})z^2,$$

where $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$ is the formal series of the δ -function.

Proof. Relation (2.13) is a trivial consequence of (2.1). Similar arguments as in the proof of Lemma (2.2) show immediately relations (2.14-2.15).

To prove the remaining relations we recall the basic property of the δ -function: for a formal series $f(z)$ with $f(a)$ defined,

$$f(z)\delta(\frac{z}{a}) = f(a)\delta(\frac{z}{a}).$$

From Proposition 2.1 it follows that for $(\alpha_i|\alpha_j) = 2$

$$[X_i^+(z), X_i^-(w)] =: X_i^+(z)X_i^-(w) : \left(\frac{z+qw}{z-qw} \frac{z+q^{-1}w}{z-q^{-1}w} - \frac{w+qz}{w-qw} \frac{w+q^{-1}z}{w-q^{-1}z} \right)$$

where the two rational functions in the parenthesis are formal series in w/z and z/w respectively.

Observe that

$$\frac{z+qw}{z-qw} \frac{z+q^{-1}w}{z-q^{-1}w} = \frac{(z+qw)(z+q^{-1}w)}{zw} \sum_{n \geq 1} \frac{q^{n+1} - q^{-n-1}}{q - q^{-1}} \left(\frac{w}{z}\right)^n$$

as an identity of formal series. Therefore we have that

$$[X_i^+(z), X_i^-(w)] =: X_i^+(z)X_i^-(w) : \frac{(z+qw)(z+q^{-1}w)}{zw(q-q^{-1})} \left(\delta(\frac{z}{w}q^{-1}) - \delta(\frac{z}{w}q) \right)$$

$$= \frac{2(q+q^{-1})}{q-q^{-1}} \left(\Psi_i(q^{-1/2}z)\delta(\frac{z}{w}q^{-1}) - \Phi_i(zq^{1/2})\delta(\frac{z}{w}q) \right),$$

where we used lemma 2.1 and the property of the δ -function.

Relation (2.18) follows from similar consideration. For $(\alpha_i|\alpha_j) = -1$ we have

$$(z - q^{\mp 1}w)(z + q^{\pm 1}w)X_i^\pm(z)X_j^\pm(w) - (q^{\mp 1}z - w)(q^{\pm 1}z + w)X_j^\pm(w)X_i^\pm(z)$$

$$=: X_i^\pm(z)X_j^\pm(w) : ((z + q^{-1}w) - (w + q^{-1}z))$$

$$= 0.$$

Similarly we proceed to show (2.19):

$$\begin{aligned}
& (z - q^{\pm 2}w)(z + q^{\mp 2}w)X_i^{\pm}(z)X_i^{\pm}(w) - (q^{\pm 2}z - w)(q^{\mp 2}z + w)X_i^{\pm}(w)X_i^{\pm}(z) \\
& =: X_i^{\pm}(z)X_i^{\pm}(w) : (z - q^2w)(z - q^{-2}w) \left(\frac{z - w}{z + w} + \frac{w - z}{w + z} \right) \\
& =: X_i^{\pm}(z)X_i^{\pm}(w) : (z - q^2w)(z - q^{-2}w)(z - w)\delta\left(-\frac{z}{w}\right) \\
& = 2(1 + q^2)(1 + q^{-2})\delta\left(-\frac{z}{w}\right)z^2
\end{aligned}$$

which completes the proof. $\square$

3. TWISTED QUANTUM ALGEBRAS

We now introduce the new quantum affine algebra in this section. Let $\mathfrak{g}$ be a complex simple Lie algebra with the standard bilinear form $(\cdot | \cdot)$ on its root lattice R as in section 2.

Definition 3.1. The algebra $U_q(\tilde{\mathfrak{g}})$ is an associative algebra generated by a_{im} , $x_{in}^{\pm}$, $\gamma = q^c$ ($n \in \mathbb{Z}, m \in 2\mathbb{Z} + 1$) with the following defining relations in terms of generating functions:

$$(3.1) \quad x_i^{\pm}(z) = \sum_{n \in \mathbb{Z}} x_{in}^{\pm} z^{-n}$$

$$(3.2) \quad \phi_i(z) = \exp((q^{-1} - q) \sum_{m > 0, \text{odd}} a_{i, -m} z^m) = \sum_{n \geq 0} \phi_{i, -n} z^n,$$

$$(3.3) \quad \psi_i(z) = \exp((q - q^{-1}) \sum_{m > 0, \text{odd}} a_{im} z^{-m}) = \sum_{n \geq 0} \psi_{in} z^{-n}.$$

The defining relation are as follows.

$$(3.4) \quad [c, \phi_i(z)] = [c, \psi_i(z)] = [c, x_i^{\pm}(z)] = 0,$$

$$(3.5) \quad [\phi_i(z), \phi_j(w)] = [\psi_i(z), \psi_j(w)] = 0,$$

$$(3.6) \quad \phi_i(z)\psi_j(w) = \psi_j(w)\phi_i(z)G_{ij}\left(\frac{z}{w}q^{-c}\right)/G_{ij}\left(\frac{z}{w}q^c\right),$$

$$(3.7) \quad \phi_i(z)x_j^{\pm}(w)\phi_i(z)^{-1} = x_j^{\pm}(w)G_{ij}(q^{\mp c/2}z/w)^{\pm 1},$$

$$(3.8) \quad \psi_i(z)x_j^{\pm}(w)\psi_i(z)^{-1} = x_j^{\pm}(w)G_{ij}(q^{\mp c/2}w/z)^{\pm 1},$$

$$(3.9) \quad [x_i^{\pm}(z), x_j^{\pm}(w)] = [x_i^{+}(z), x_j^{-}(w)] = 0, \quad (\alpha_i | \alpha_j) = 0$$

$$(3.10) \quad (z + w)[x_i^{+}(z), x_j^{-}(w)] = 0, \quad (\alpha_i | \alpha_j) = -1$$

$$\begin{aligned}
(3.11) \quad [x_i^{+}(z), x_i^{-}(w)] &= \frac{2(q + q^{-1})}{q - q^{-1}} \left(\psi_i(q^{-c/2}z)\delta\left(\frac{w}{z}q^c\right) - \phi_i(zq^{c/2})\delta\left(\frac{w}{z}q^{-c}\right) \right), \\
& (z - q^{\mp 1}w)(z + q^{\pm 1}w)x_i^{\pm}(z)x_j^{\pm}(w) = \\
& (q^{\mp 1}z - w)(q^{\pm 1}z + w)x_j^{\pm}(w)x_i^{\pm}(z), \quad (\alpha_i | \alpha_j) = -1,
\end{aligned}$$

$$(3.12) \quad (z - q^{\pm 2}w)(z + q^{\mp 2}w)x_i^{\pm}(z)x_i^{\pm}(w) = (q^{\pm 2}z - w)(q^{\mp 2}z + w)x_i^{\pm}(w)x_i^{\pm}(z)$$

$$(3.13) \quad + 2(q + q^{-1})^2 \delta\left(-\frac{w}{z}\right)c.$$

where G_{ij} is defined in (2.12).

In the Drinfeld realization [Dr] of the usual quantum affine algebras there are several relations similar to our relations, but with a different definition of $G_{ij}(z)$. Another obvious difference is that we have more complicated commutation relations between operators $x_i^\pm(z)$ of the same type.

The twisted algebra $U_q(\tilde{\mathfrak{g}})$ has a Hopf algebra structure. Let $c_1 = c \otimes 1$ and $c_2 = 1 \otimes c$.

$$(3.14) \quad \Delta(x_i^+(z)) = x_i^+(z) \otimes 1 + \phi_i(q^{c_1/2}z) \otimes x_i^+(q^{c_1}z),$$

$$(3.15) \quad \Delta(x_i^-(z)) = 1 \otimes x_i^-(z) + x_i^-(q^{c_2}z) \otimes \phi_i(q^{c_2/2}z),$$

$$(3.16) \quad \Delta(\phi_i(z)) = \phi_i(q^{-c_2/2}z) \otimes \phi_i(q^{c_1/2}z),$$

$$(3.17) \quad \Delta(\psi_i(z)) = \psi_i(q^{c_2/2}z) \otimes \psi_i(q^{-c_1/2}z),$$

$$(3.18) \quad \Delta(c) = c \otimes 1 + 1 \otimes c.$$

The antipode is given by:

$$(3.19) \quad S(x_i^+(z)) = -\phi_i(q^{-c/2}z)x_i^+(q^{-c}z),$$

$$(3.20) \quad S(x_i^-(z)) = -x_i^-(q^{-c}z)\psi_i(q^{-c/2}z)^{-1},$$

$$(3.21) \quad S(\phi_i(z)) = \phi_i(z)^{-1},$$

$$(3.22) \quad S(\psi_i(z)) = \psi_i(z)^{-1},$$

$$(3.23) \quad S(c) = -c.$$

The counit map is

$$(3.24) \quad \epsilon(x_i^\pm(z)) = 0, \quad \epsilon(\phi_i(z)) = 1, \quad \epsilon(\psi_i(z)) = 1$$

$$(3.25) \quad \epsilon(q^c) = 1.$$

The relations (2.17-2.19) resolve the Serre relations for our algebra. We remark that our algebra is different from the algebra of Ding-Iohara [DI] due to our relations (2.17-2.19). If we replace the relation (2.19) by the following weaker one:

$$\begin{aligned} & (z+w)(z-q^{\pm 2}w)(z+q^{\mp 2}w)X_i^\pm(z)X_i^\pm(w) \\ &= (z+w)(q^{\pm 2}z-w)(q^{\mp 2}z+w)X_i^\pm(w)X_i^\pm(z). \end{aligned}$$

Then this modified algebra has a common quotient with the Ding-Iohara algebra in [DI].

4. A TWISTED QUANTUM KAC-MOODY ALGEBRA AND ITS LEVEL ONE REPRESENTATIONS

We can generalize the algebra to any even lattice. Let Q be an arbitrary integral lattice generated by α_i , $i = 1, \dots, l$, equipped with a symmetric form $(\cdot | \cdot)$ such that

$$(4.1) \quad (\alpha_i | \alpha_j) \in \mathbb{Z}, \quad (\alpha_i | \alpha_i) = 2.$$

In the following we can view this lattice as that of a root lattice of Kac-Moody algebra [K1]. In this sense the following construction parallels to the quantum loop algebra associated to the Kac-Moody algebra studied in [J7].

Let $U_q(\tilde{h}_Q)$ be the associative algebra generated by $a_i(m)$, $m \in \mathbb{Z}$ and $\gamma^{\pm 1}$ with the defining relation (2.1-2.2). Then we can consider the central extension of Q by

a finite group $\langle \kappa \rangle$ similarly (cf. [FLM]). If we use a_i to denote the preimage of α_i , then we have

$$(4.2) \quad a_i a_j = (-1)^{(\alpha_i | \alpha_j)} a_j a_i.$$

Let T be any $\hat{Q}$ -module such that $\kappa = \omega$, a $o(\kappa)$ th primitive root of unity. The vertex space is defined accordingly:

$$(4.3) \quad V_Q = S(\tilde{h}_Q^-) \otimes T.$$

For each α_i we define the vertex operator by the same expression as in (2.7). To compute the operator product expansion we need the following q -analogs of series $(z - w)^r$ introduced in [J5, J7]. For $r \in \mathbb{C}$ we define the q -analog to be in base q^2 :

$$(4.4) \quad (1 - z)_{q^2}^r = \frac{(q^{-r+1}z; q^2)_\infty}{(q^{r+1}z; q^2)_\infty} = \exp\left(-\sum_{n=1}^{\infty} \frac{[rn]}{n[n]} z^n\right),$$

where $(b; q)_\infty = \prod_{n=0}^{\infty} (1 - q^n b)$. The twisted q -analog is defined by

$$(4.5) \quad \left(\frac{1 - z}{1 + z}\right)_{q^2}^r = \frac{(1 - z)_{q^2}^r}{(1 + z)_{q^2}^r} = \exp\left(-\sum_{n \in \mathbb{Z}_1^+} \frac{2[rn]}{n[n]} z^n\right),$$

The following OPE can be calculated similarly as in section 2.

Proposition 4.1.

$$\begin{aligned} X_i^\pm(z) X_j^\pm(w) &=: X_i^\pm(z) X_j^\pm(w) : \left(\frac{1 - q^{\mp 1} w/z}{1 + q^{\mp 1} w/z}\right)_{q^2}^{(\alpha_i | \alpha_j)}, \\ X_i^\pm(z) X_j^\mp(w) &=: X_i^\pm(z) X_j^\mp(w) : \left(\frac{1 + w/z}{1 - w/z}\right)_{q^2}^{(\alpha_i | \alpha_j)}. \end{aligned}$$

Other commutations on the vertex representation space V_Q are given as follows.

Theorem 4.1. *The twisted vertex operators satisfy the following relations:*

$$(4.6) \quad [\Phi_i(z), \Psi_j(w)] = [\Phi_i(z), \Psi_j(w)] = 0,$$

$$(4.7) \quad \Phi_i(z) X_j^\pm(w) \Phi_i(z)^{-1} = X_j^\pm(w) G_{ij}(q^{\mp 1/2} z/w)^{\pm 1},$$

$$(4.8) \quad \Psi_i(z) X_j^\pm(w) \Psi_i(z)^{-1} = X_j^\pm(w) G_{ij}(q^{\mp 1/2} w/z)^{\pm 1},$$

$$(4.9) \quad [X_i^\pm(z), X_j^\pm(w)] = [X_i^+(z), X_j^-(w)] = 0, \quad (\alpha_i | \alpha_j) = 0,$$

$$(4.10) \quad (z + w)_{q^2}^{-(\alpha_i | \alpha_j)} [X_i^+(z), X_j^-(w)] = 0, \quad (\alpha_i | \alpha_j) < 0$$

$$[X_i^+(z), X_i^-(w)] = \frac{2(q + q^{-1})}{q - q^{-1}} \left(\Psi_i(q^{-1/2} z) \delta\left(\frac{w}{z} q\right) - \Phi_i(z q^{1/2}) \delta\left(\frac{w}{z} q^{-1}\right) \right),$$

$$(4.11) \quad (z - w)_{q^2}^{-(\alpha_i | \alpha_j) - 1} (z - q^{(\alpha_i | \alpha_j)} w) (z + q^{-(\alpha_i | \alpha_j)} w) X_i^\pm(z) X_j^\pm(w) =$$

$$(z - w)_{q^2}^{-(\alpha_i | \alpha_j) - 1} (q^{(\alpha_i | \alpha_j)} z - w) (q^{-(\alpha_i | \alpha_j)} z + w) X_j^\pm(w) X_i^\pm(z), \quad (\alpha_i | \alpha_j) < 0,$$

$$(4.12) \quad (z - q^{\pm 2} w) (z + q^{\mp 2} w) X_i^\pm(z) X_i^\pm(w) = (q^{\pm 2} z - w) (q^{\mp 2} z + w) X_i^\pm(w) X_i^\pm(z)$$

$$(4.13) \quad + 2(q + q^{-1})^2 \delta\left(-\frac{w}{z}\right).$$

Remark 4.2. Note that we use the factor $(z+w)_{q^2}^{-(\alpha_i|\alpha_j)}$ and $(z-w)_{q^2}^{-(\alpha_i|\alpha_j)-1}$ to suppress poles in (4.10) and (4.12) respectively. We can cancel these two polynomials and rewrite the commutation relations in terms of delta functions in a representation. It is always possible to refine the commutation relations as in the classical case [K2] and quantum case [J7]. In the quantum case one uses the q -difference of the delta function $\delta(z-w)$ to expand zero expressions. As in [J7] we have for $n \geq 0$

$$(4.14) \quad i_{z,w}(z-w)_{q^2}^{-n-1} - i_{w,z}(z-w)_{q^2}^{-n-1} = \partial_{q,w}^{(n)} \delta(z-w),$$

where $i_{z,w}$ means the expansion in the range $|z| \gg |w|$ and the $\partial_{q,w}^{(n)}$ denote the q -divided powers of n th q -difference ∂_q :

$$(4.15) \quad \partial_q f(z) = \frac{f(qz) - f(q^{-1}z)}{(q - q^{-1})z}.$$

The delta function $\delta(z-w)$ means $z\delta(z/w)$. Then we can rewrite for example the relation (4.11)

$$\begin{aligned} & (z - q^{(\alpha_i|\alpha_j)}w)(z + q^{-(\alpha_i|\alpha_j)}w)X_i^\pm(z)X_j^\pm(w) \\ & - (q^{(\alpha_i|\alpha_j)}z - w)(q^{-(\alpha_i|\alpha_i)}z + w)X_j^\pm(w)X_i^\pm(z) \\ & =: X_i^\pm(z)X_j^\pm(w) : (z+w)_{q^2}^{-(\alpha_i|\alpha_j)-1} \partial_{q,w}^{(-(\alpha_i|\alpha_j))} \delta(z-w), \end{aligned}$$

where $(\alpha_i|\alpha_j) < 0$.

Definition 4.3. Let $\mathfrak{g}$ be a Kac-Moody algebra with the root lattice $Q = \mathbb{Z}\alpha_1 \oplus \cdots \oplus \mathbb{Z}\alpha_l$ and the symmetric bilinear form $(\cdot | \cdot)$. The algebra $U_q(\widehat{\mathfrak{g}})$ is an associative algebra with generators $a_{im}, x_{in}^\pm, \gamma = q^c$ ($n \in \mathbb{Z}, m \in 2\mathbb{Z}+1$) and following defining relations:

$$(4.16) \quad [c, \Phi_i(z)] = [c, \Psi_i(w)] = [c, X_i^\pm(z)] = 0$$

$$(4.17) \quad [\Phi_i(z), \Psi_j(w)] = [\Phi_i(z), \Psi_j(w)] = 0,$$

$$(4.18) \quad \Phi_i(z)X_j^\pm(w)\Phi_i(z)^{-1} = X_j^\pm(w)G_{ij}(q^{\mp c/2}z/w)^{\pm 1},$$

$$(4.19) \quad \Psi_i(z)X_j^\pm(w)\Psi_i(z)^{-1} = X_j^\pm(w)G_{ij}(q^{\mp c/2}w/z)^{\pm 1},$$

$$(4.20) \quad [X_i^\pm(z), X_j^\pm(w)] = [X_i^+(z), X_j^-(w)] = 0, \quad (\alpha_i|\alpha_j) = 0,$$

$$(4.21) \quad (z+w)_{q^2}^{-(\alpha_i|\alpha_j)}[X_i^+(z), X_j^-(w)] = 0, \quad (\alpha_i|\alpha_j) < 0$$

$$[X_i^+(z), X_i^-(w)] = \frac{2(q+q^{-1})}{q-q^{-1}} \left(\Psi_i(q^{-c/2}z)\delta\left(\frac{w}{z}q^c\right) - \Phi_i(zq^{c/2})\delta\left(\frac{w}{z}q^{-c}\right) \right),$$

$$(4.22) \quad (z-w)_{q^2}^{-(\alpha_i|\alpha_j)-1}(z-q^{(\alpha_i|\alpha_j)}w)(z+q^{-(\alpha_i|\alpha_j)}w)X_i^\pm(z)X_j^\pm(w) = \\ (z-w)_{q^2}^{-(\alpha_i|\alpha_j)-1}(q^{(\alpha_i|\alpha_j)}z-w)(q^{-(\alpha_i|\alpha_i)}z+w)X_j^\pm(w)X_i^\pm(z), \quad (\alpha_i|\alpha_j) < 0,$$

$$(4.23) \quad (z-q^{\pm 2}w)(z+q^{\mp 2}w)X_i^\pm(z)X_i^\pm(w) = (q^{\pm 2}z-w)(q^{\mp 2}z+w)X_i^\pm(w)X_i^\pm(z) \\ + 2(q+q^{-1})^2\delta\left(-\frac{w}{z}\right)c.$$

$$(4.24)$$

where the generating functions are defined by (3.2-3.3).

We close this section with the following statement.

Conjecture 4.1. *The quantum affine algebra $U_q(\hat{sl}_2)$ is isomorphic to the twisted algebra generated by $a_{1m}(m \in 2\mathbb{Z} + 1), \gamma, x_{1n}^\pm(n \in \mathbb{Z})$ subject to the relations (3.4-3.13) associated to A_1 .*

At $c = 1$ this statement is true by specializing $q \rightarrow 1$ [LW], however we do not know the exact isomorphism at the quantum level. The exact isomorphism will solve the so-called principal vertex construction of $U_q(\hat{sl}_2)$.

REFERENCES

- [DF] J. Ding and B. Feigin, Quantum current operators–II. Difference equations and quantum parafermion construction, Publ. RIMS Kyoto University **33** (1997), 285–300.
- [DI] J. Ding and K. Iohara, Generalization and deformation of Drinfeld quantum affine algebras, Lett. Math. Phys. **41** (1997), 181–193.
- [DL] C. Dong and J. Lepowsky, The algebraic structure of relative twisted vertex operators, J. Pure & Applied Algebra **110** (1996), 259–295.
- [Dr] V.G. Drinfeld, A new realization of Yangians and quantized affine algebras, Soviet Math. Dokl. **36** (1988), 212–216.
- [FJ] I.B. Frenkel and N. Jing, Vertex representations of quantum affine algebras, Proc. Natl. Acad. Sci. USA **85** (1988), 9373–9377.
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex operator algebras and the monster*, Academic Press, New York, 1988.
- [Jb] M. Jimbo, A q -difference analog of $U(\mathfrak{g})$ and the Yang-Baxter equation, Lett. Math. Phys. **10** (1985), 63–69.
- [J1] N. Jing, Vertex operators, symmetric functions and their q -deformations, dissertation, Yale Univ., 1989.
- [J2] N. Jing, Twisted vertex representations of quantum affine algebras, Invent. Math. **102** (1990), 663–690.
- [J3] N. Jing, Vertex operators, symmetric functions, and the spin group Γ_n , J. Algebra **138** (1991) 340–398.
- [J4] N. Jing, Vertex operators and generalized symmetric functions, in: Proc. of Conf. on Quantum Topology (KSU, March 1993), ed. D. Yetter, World Scientific, Singapore, 1994, pp. 111–126.
- [J5] N. Jing, Higher level representations of the quantum affine algebra $U_q(\hat{sl}(2))$, J. Algebra **182** (1996), 448–468.
- [J6] N. Jing, Quantum Z -algebras and representations of quantum affine algebras, Commun. Algebra, to appear. (math.QA/9806043)
- [J7] N. Jing, Vertex representations of the quantum Kac-Moody algebras, Lett. Math. Phys. **44** (1998), no. 4, 261–271.
- [K1] V.G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.
- [K2] V.G. Kac, *Vertex algebras for beginners*, Univ. Lect. Ser. **10**, Amer. Math. Soc., Providence, RI, 1997.
- [L] J. Lepowsky, Calculus of twisted vertex operators, Proc. Natl. Acad. Sci. USA **82** (1985), 8295–8299.
- [LP] J. Lepowsky and P. Primc, Structure of the standard modules for affine Lie algebra $A_1^{(1)}$, Contemp. Math. **46**, Amer. Math. Soc., Providence, RI, 1985.
- [LW] J. Lepowsky and R.L. Wilson, Construction of the affine Lie algebra $A_1^{(1)}$, Commun. Math. Phys. **62** (1978), 43–53.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695-8205, U.S.A.

E-mail address: jing@math.ncsu.edu